

18 Robert W. Klaver, Earth Resources Observation and Science Center, U.S. Geological
19 Survey, 47914 252nd Street, Sioux Falls, South Dakota, 57198, USA. Email:
20 bklaiver@usgs.gov, phone: 605-594-6067.

21 *Abstract:* Greater sage-grouse (*Centrocercus urophasianus*) populations in North Dakota
22 declined approximately 67% between 1965 and 2003 and are listed as a Priority
23 Level 1 Species of Special Concern by the North Dakota Game and Fish
24 Department. The habitat and ecology of this species at the eastern edge of its
25 historical range is largely unknown. We investigated nest site selection by greater
26 sage-grouse and nest survival in North Dakota during 2005 and 2006. Sage-grouse
27 selected nest sites in sagebrush (*Artemisia* spp.) with more total vegetative cover,
28 greater sagebrush density, and greater 1-m visual obstruction from the nest than
29 random sites. Height of grass and shrub (sagebrush) at nest sites were shorter than
30 random sites because areas where sagebrush was common were sites in low seral
31 condition or dense clay or clay-pan soils with low productivity. Constant survival
32 estimates of incubated nests were 33% in 2005 and 30% in 2006. Variables that
33 described the resource selection function for nests were not those that modeled nest
34 survival. Nest survival was positively influenced by percent shrub (sagebrush)
35 cover and grass height. Daily nest survival decreased substantially when percent
36 shrub cover declined below about 9% and when grass heights were less than about
37 16 cm. Daily nest survival rates decreased with increased daily precipitation.

38 **Introduction**

39 Greater sage-grouse (*Centrocercus urophasianus*) populations have declined by
40 45% to 80% across their range (Schroeder et al. 2004, Connelly et al. 2004). In North
41 Dakota, greater sage-grouse (hereafter sage-grouse) declined approximately 67% from
42 1965 to 2003 (Connelly et al. 2004). While sage-grouse in North Dakota are genetically
43 contiguous with populations in Montana and South Dakota (Oyler-McCance et al. 2005),
44 they could become isolated by conversion of sagebrush to agriculture (Smith et al. 2005),
45 and oil and gas development (Connelly et al. 2004, Walker et al. 2007). Long-term
46 population declines in sage-grouse in North Dakota (Smith et al. 2004) have resulted in
47 classifying sage-grouse as Priority Level 1 Species of Special Concern in North Dakota
48 (McCarthy & Kobriger 2005). Altered habitat quality and quantity in sagebrush
49 ecosystems (e.g., Connelly et al. 2004, Welch 2005) may result in low survival and
50 productivity in sage-grouse (Aldridge & Brigham 2001) and declining populations.
51 Therefore, understanding characteristics important to selection of nest sites and factors that
52 affect nest survival is critical to the management, conservation, and rehabilitation of
53 sagebrush habitats for sage-grouse. Despite well understood reproductive ecology in the
54 core of sage-grouse range, knowledge of reproductive ecology and habitat selection by
55 sage-grouse occurring at the eastern edge of their distribution is limited. Therefore, our
56 objective was to quantify nest habitat selection by sage-grouse in North Dakota and
57 determine specific factors associated with survival of sage-grouse nests. These data will
58 help in the development of management recommendations to assist state and federal
59 agencies in managing habitats for sage-grouse.

60 **Materials and methods**

61 *Study area*

62 Our study area included Bowman and Slope counties in southwestern North Dakota
63 and Fallon County in southeastern Montana. The area has flat to gently-rolling prairie,
64 with a few buttes and intermittent streams. Annual precipitation ranges from 35.6 cm to
65 40.6 cm, most of which occurs from April to September. Summer temperatures range
66 from 9.9°C to 27.5°C and winter temperatures range from -15.6°C to 0.2°C (North Dakota
67 Agricultural Weather Network, 2006).

68 Vegetation communities included mixed grass prairie with perennial and annual
69 forbs and grasses and shrub steppe comprised of silver sagebrush (*Artemisia cana*),
70 Wyoming big sagebrush (*A. tridentata wyomingensis*), rubber rabbitbrush (*Chrysothamnus*
71 *nauseosus*), and greasewood (*Sarcobatus vermiculatus*). Common grasses included
72 western wheatgrass (*Pascopyrum smithii*), Kentucky bluegrass (*Poa pratensis*), Japanese
73 brome (*Bromus japonicus*), needle and thread (*Stipa comata*), and junegrass (*Koeleria*
74 *macrantha*). Forbs included common dandelion (*Taraxacum officinale*), common yarrow
75 (*Achillea millefolium*), and textile onion (*Allium textile*) (Johnson & Larson 1999).

76 Most of the land is privately owned and the primary land use is cattle ranching.
77 Areas managed by the Bureau of Land Management (BLM) for grazing were stocked at 4
78 -10 acres per animal unit month (AUM) under continuous or rotational grazing that begins
79 in early to mid June (Mitch Iverson, Belle Fourche BLM, personal communications). Oil
80 and gas development is extensive in some areas.

81 *Capture and telemetry* -- We captured female sage-grouse at night on or near leks
82 from late March through April in 2005 and 2006 (Wakkinen et al. 1992b). We recorded
83 age (Crunden 1963) and placed a 20-gram necklace radio transmitter (Advanced Telemetry
84 Systems, Isanti, Minnesota) with mortality sensors on each female. The transmitters were
85 less than 2% of the body weight of individually marked sage-grouse. All field methods
86 complied with the Institutional Animal Care and Use Committee (07-A032) at South
87 Dakota State University.

88 We located females 2-3 times each week from the time they were captured and
89 radiomarked, until the outcome of nesting had been determined aided with a hand-held
90 3-element yagi antenna. Nests were inconspicuously marked with plastic flagging >20 m
91 south of the nest, near or at ground level to avoid making them visible except on close
92 inspection. The nest location was recorded with a GPS. Occasionally, the interval
93 between telemetry locations was greater than 2-3 days because weather prohibited access
94 to the nest area. Therefore, onset of incubation could not be accurately estimated from
95 behavior of females. As a result, we flushed incubating hens and estimated nest initiation
96 by backdating from incubation stage estimated from egg flotation (Hays & LeCroy 1971)
97 and adding 1.3 days for each egg laid (Patterson 1952). If the female was absent from the
98 nest area for more than 3 consecutive locations, we approached the nest site to determine
99 fate. Success or failure of nests was determined by membrane conditions of the eggs
100 (Klebenow 1969) or observation of a brood with the radiocollared female. Nests were
101 considered successful if ≥ 1 egg hatched.

102 *Vegetation measurements* -- During May and June each year of the study, we
103 characterized vegetation at nest sites and random sites. Most sage-grouse nest within 3.2
104 km of a lek (Braun et al. 1977, Aldridge & Brigham 2001) so we selected random sites
105 from which to estimate resource selection for nesting from within a 3-km buffer
106 surrounding leks on which we observed sage-grouse. Because nest sites are normally
107 located beneath sagebrush (Connelly et al. 2000), random sites were selected at the nearest
108 sagebrush plant to the random coordinates. We recorded slope and aspect at each nest and
109 random site using a clinometer and compass.

110 We established 4 50-m transects that were centered over the nest or random site.
111 We recorded species, height, length, and width of sagebrush plants at the intersection of
112 these transects. Using 10-m intervals ($n = 20$) along each transect we recorded the distance
113 to the nearest shrub (usually sagebrush) using a point-centered-quarter method (Cottam &
114 Curtis 1956), and recorded the species, height, length, and width of each shrub. We also
115 recorded the maximum height of grass growing from beneath the sagebrush. We estimated
116 visual obstruction (VOR) and height of grass at each nest site, and for each meter out to
117 5 m from the nest, and then at 10-m intervals along each transect using a modified Robel
118 pole delineated in 2.54-cm increments (Robel et al. 1970, Benkobi et al. 2000). We used
119 the Daubenmire (1959) method to estimate canopy cover of vegetation. This method is
120 amenable to collecting data on windy days and yields data that are similar (<3% difference
121 for sagebrush) to the line-intercept method (Floyd & Anderson 1987), and may provide
122 more accurate estimates than line-intercept methods (Booth et al. 2006). We estimated
123 canopy cover from a height of about 1 m in 24, 0.1-m² quadrats (Daubenmire 1959). Four

124 quadrats were placed at the intersection of the transects (over the nest) and at the four
125 terminal ends of 1-m legs forming the pattern of an H every 10 m along each transect. We
126 estimated percent canopy cover for total cover, total shrubs, total forbs, total grasses, litter,
127 bare ground, sagebrush and dominant species of grasses and forbs using six categories
128 (Daubenmire 1959). We obtained measures of maximum and minimum daily temperature
129 and daily precipitation throughout the nesting season from the closest weather station in
130 Bowman County (North Dakota Agricultural Weather Network 2006).

131 *Data analyses*

132 *Nesting* -- We tested for differences in clutch size distributions between adults and
133 yearlings using chi-square goodness of fit test. Chi-square goodness of fit tests were used
134 to test differences in nest initiation rates between years and among ages of females. We
135 calculated distance from each nest to the center of nearest lek and distance from each nest
136 to lek of capture (if the hen was captured that year) using corresponding GPS coordinates.
137 We tested for differences in these distributions between successful and unsuccessful nests,
138 and between adults and yearling hens using multiple response permutation program
139 (MRPP; Mielke & Berry 2001). Statistical significance was determined at $\alpha \leq 0.05$ for
140 these tests.

141 *Habitat selection* -- Average percentage canopy cover was recorded for each
142 variable at nests and random sites. Visual obstruction was calculated at the nest and at 1 m
143 intervals out to 5 m. Average VOR was also calculated for each site. We estimated
144 sagebrush density using maximum likelihood estimates of point-centered-quarter method
145 (Pollard 1971). We then used MRPP to test the distribution of vegetation characteristics

146 between nests and random sites to distinguish important variables to include in models of
147 nest survival and selection of nest sites. Statistical significance was determined with a
148 critical value of $\alpha \leq 0.05$.

149 We used an information theoretic approach (Burnham & Anderson 2002) with
150 logistic regression to estimate models depicting vegetation characteristics selected by
151 female sage-grouse for nests. Because we had a very large number of variables from
152 estimated canopy cover by species and collected extensive measurements, we developed
153 10 candidate models that included variables that exhibited differences between nest and
154 random sites from MRPP tests (see Hosmer & Lemeshow 2000, Guthery et al. 2005,
155 Stephens et al. 2005). These models included percent total cover, percent grass cover,
156 percent forb cover, percent sagebrush cover, sagebrush height, site-VOR, nest VOR, 1-m
157 VOR, grass height from the Robel pole, and sagebrush density. Year was considered as a
158 design variable and was included in all candidate models. We tested the strength of the
159 best predictive model of nest sites selected using receiver operating characteristic curves
160 (ROC). Receiver operating characteristic curve values between 0.8 and 0.9 were
161 considered excellent discrimination, and ROC values between 0.7 and 0.8 were considered
162 acceptable discrimination (Hosmer & Lemeshow 2000). The statistical tests described
163 above were made using SPSS (2002) or SAS (2005).

164 *Nest survival* -- We estimated daily survival rate (DSR) of nests using the nest
165 survival model in Program MARK (White & Burnham 1999, Dinsmore et al. 2002). We
166 established 6 May as first nest day.

167 Nest survival probabilities were estimated as a function of age of hen, nest age, and
168 vegetation characteristics at nests. We then modeled effects of time-dependent variables
169 year, maximum and minimum daily temperature, and daily precipitation using the best
170 survival models from the previous analysis also using program MARK (White & Burnham
171 1999). Continuous covariates were standardized as deviations from a mean of 0.
172 Categorical and time-dependent covariates were coded with the actual values so they
173 would not hamper numerical optimization of likelihood (Burnham & Anderson 2002).

174 **Results**

175 Thirty-nine hens were captured and fitted with necklace-mounted radio transmitters
176 during spring 2005 and 2006 (21 during 2005, 18 during 2006); 36% were adults. Eleven
177 females in 2006 were marked in 2005.

178 *Nesting*

179 Adults initiated nests approximately 5 days earlier than yearlings. Nests were 6 to
180 8 days on average into incubation when detected. There were 2 renests in 2005 that were
181 initiated in mid to late May; no renesting occurred in 2006. Renesting rate was 10%. All
182 radiocollared hens initiated a nest in 2005. In 2006, 13 of 14 adults (93%), and 5 of 7
183 yearlings (71%) incubated a nest (includes those that abandoned). There was no difference
184 in nest initiation rates between years ($P = 0.11$). Nest initiation rate (including those that
185 abandoned) for adult hens ($n = 20$) was 95% and did not differ ($P = 0.58$) from yearling
186 hens (88%, $n = 16$). Nest initiation averaged 92% across age groups and years.

187 For nests that we could determine clutch size, ($n = 33$) average clutch size was
188 7.9 ± 0.5 eggs. There was no difference in clutch size between adults and yearlings

189 ($P = 0.86$). We eliminated four nests from further analyses because we believed they were
190 abandoned because of disturbance from our field crews. In 2005, 3 of the nests were
191 abandoned by the hen, and 5 were depredated. In 2006, 1 nest was abandoned by the hen,
192 and 8 nests were depredated.

193 The average distance from nests to the lek at which a hen was captured was
194 4.9 ± 4.1 ($\bar{x} \pm \text{SE}$) km, and did not differ ($P = 0.67$) between successful and unsuccessful
195 nests. The average distance from nests to the nearest lek was 2.7 ± 2.4 km. Unsuccessful
196 nests did not differ from successful nests in relation to distance to nearest lek ($P = 0.45$).
197 Average distance to the nearest lek did not differ ($P = 0.45$) between years nor ($P = 0.77$)
198 between adults or yearlings.

199 *Nest selection*

200 Eighty-five percent of nests were located under Wyoming big sagebrush ($n=29/34$).
201 Other than sagebrush, one nest each was located beneath four-wing saltbush (*Atriplex*
202 *canesens*), eastern redcedar (*Juniperus virginiana*), and wheat stubble (*Triticum* spp.), and
203 two were in sweet clover. Vegetation at random sites was sparse, but slightly taller than
204 nest sites. Sage-grouse nest sites had greater ($P \leq 0.05$) percent canopy cover of total
205 vegetation (total cover), grass cover, forb cover, sagebrush cover, and litter (Table 1).
206 Moreover, nest sites had greater visual obstruction at the nest (nest VOR) and surprisingly
207 even greater visual obstruction 1 m away (1-m VOR). Although vegetation was taller at
208 random sites, VOR for nest sites was greater ($P < 0.01$) than random sites. Sagebrush
209 density also was greater ($P < 0.01$) at nest sites than random sites (Table 1). Intermediate
210 wheatgrass (*Thinopyrum intermedium*) was the only dominant grass with greater canopy at

211 random sites; otherwise canopy cover of dominant grasses was greater at nest sites.
212 Nonetheless grass height and shrub height were marginally taller ($P > 0.07$) at random
213 sites.

214 We included models with total cover, shrub density, shrub height, grass height, nest
215 VOR and 1-m VOR in the evaluation of nest resource selection. Other variables were
216 excluded because of correlations with these variables. Because total cover exhibited the
217 smallest individual variable deviance, we constructed iterations of models around this
218 variable. Of the 25 models we considered, 5 models were included in the set with ΔAIC_c
219 < 2 (Table 2). Two models, both including total cover, and shrub height, with nest VOR
220 (highest rank), and 1-m VOR (second highest rank) best explained the nest resource
221 selection by female sage-grouse. Nest sites were positively associated with greater
222 percent total cover, greater 1-m VOR and nest VOR, and negatively related to shrub
223 height. The third ranked model included only total cover and shrub height. Models that
224 also included grass height, although ranked in the “supported” set, did not improve the
225 deviance suggesting that grass height was not really important in describing the resource
226 selection by nesting sage-grouse. The odds ratios indicated that nest VOR was the most
227 important of the variables in the model. An increment of 2.54 cm for nest VOR increased
228 the predicted probability of the site to be a nest by $16\% \pm 1\%$ (95%CI). Increasing total
229 vegetative cover by 10% increased the predicted probability of the site being selected for
230 nesting by a $0.60\% \pm 0.3\%$ (CI 95%). Finally the odds ratio for shrub height indicated a
231 $9.1\% \pm 1.3\%$ decreased the predicted probability of a nest with each 1 cm increase in shrub
232 height. Classification accuracy of the model was acceptable with an ROC value = 0.82.

233 Odds ratios for the second ranked model were virtually identical to the previous model,
234 except that greater weight was placed on the 1-m VOR.

235 *Nest survival*

236 Fourteen hens were included in nest survival analyses in 2005 (8 yearlings, 6
237 adults), and 15 hens were included in nest survival analyses in 2006 (3 yearlings, 12
238 adults). Nest survival did not differ between years ($P < 0.05$). Estimated constant nest
239 survival was 33% in 2005 ($n = 14$) and 30% in 2006 ($n = 15$).

240 The best model from the nest site selection (above) was the lowest ranked model
241 describing nest survival of the 41 models considered (Table 3). There was virtually no
242 support for any of the single variable models. We included precipitation, constant
243 survival, nest age and year in the table because they are often variables interest despite
244 their lack of support in our models. These were the same three top models when we
245 included only vegetation characteristics (minus precipitation). Before including
246 precipitation, AIC_w for these models were 0.15, 0.14, and 0.06, respectively. Although
247 precipitation by itself provided little insight into nest survival of sage-grouse, when include
248 with vegetation characteristics, the three top vegetation models showed strong support,
249 with the fourth best model including just vegetation.

250 Evaluation of the coefficients of the second ranked model suggested that adding
251 total cover to the model brought in some strong variable interactions causing the intercept
252 to be below zero. Therefore, we eliminated this model and AIC_w assigned to models
253 without this are in parentheses. Unequivocally, the model with the greatest support was
254 the model including grass height, shrub cover, nest VOR and precipitation. The addition

255 of site VOR to the highest ranked model contributed very little reducing the deviance in
256 the model. Therefore, we were left with highest ranked model to interpret.

257 Survival of sage-grouse nests was positively associated with grass height and shrub
258 cover, and negatively associated with precipitation, and nest VOR. The relation between
259 nest VOR and survival seemed counter intuitive, so we examined measurements of the nest
260 shrub and found that successful nests were indeed in shorter shrubs than unsuccessful
261 nests. None of the 95% CI's for odds ratios included 0. Grass height and shrub cover
262 increasing the probability of nests surviving by about 1.2% for each unit increase. An
263 increase in nest VOR of 2.5 cm decreased the chances of the nest surviving by 2% and 1
264 cm of precipitation decreased the chances of a nest surviving by about 7%. Daily
265 precipitation and had a consistent negative effect on nest survival (Figure 1), which was
266 amplified when shrub cover was less than about 9% or when grass height was less than
267 about 16 cm (Figure 1C).

268 **Discussion**

269 *Breeding chronology and nesting*

270 Average clutch size in southwestern North Dakota was similar to that throughout
271 the range of sage-grouse (Wallestad & Pyrah 1974, Sveum 1995). Despite predictions of
272 age-specific differences in clutch size (Wallestad & Pyrah 1974, Petersen 1980), adults and
273 yearlings laid similar clutch sizes in our study.

274 We interpret earlier nesting by adults to their being physiologically more mature
275 and ready for reproduction than yearlings (e.g., Schroeder 1997). Improved habitat
276 (nutritional) quality was postulated to be responsible increased production in sage-grouse

277 in Oregon (Barnett & Crawford 1994), and Gregg et al. (2006) showed that hens with
278 greater plasma protein were more likely to re-nest. The low re-nesting rate in our study
279 suggests that some aspect of the habitat was lacking.

280 Sage-grouse do not always nest near a lek, and may nest independent of lek
281 locations (Bradbury et al. 1989, Wakkinen et al. 1992a). In Alberta Canada, less than ½
282 (41%) of nests were within 3.2 km of the lek (Aldridge & Brigham 2001). However
283 elsewhere, most nests occur within 3.2 km of leks (Braun et al. 1977). The population of
284 sage-grouse we studied was non-migratory and 68% of nests were within 3.2 km of a lek;
285 86% of nests were within 5 km of a lek. It is likely that suitable nesting habitat for sage-
286 grouse in the Dakotas, at the eastern fringe of sage-grouse range, occurs near leks.
287 Aldridge & Brigham (2001) came to a similar conclusion for a population at the northern
288 fringe of the sage-grouse range. While in more contiguous sagebrush of Wyoming, there
289 was less propensity for sage-grouse to nest near the lek (Holloran & Anderson 2005).

290 *Nest selection*

291 Most studies describe the importance of sagebrush canopy cover and herbaceous
292 canopy cover (Wakkinen 1990, Connelly et al. 1991, Sveum et al. 1998, Hagen et al. 2007)
293 for sage-grouse nesting habitat. However in southwestern North Dakota, nest resource
294 selection may take on different characteristics than other portions of the sage-grouse range.
295 Sage-grouse usually nest in taller sagebrush (Connelly et al. 2000) and often select the
296 tallest sagebrush (Wakkinen 1990, Apa 1998). However in our study, sage-grouse selected
297 sites with greater vegetative cover and greater visual obstruction at and near the nest, but
298 with shorter shrub height than was available in the area. Tall grass also can be important

299 contributor to concealment of sage-grouse nests (Connelly et al. 1991, Gregg et al. 1994)
300 and although grass height was included in models with marginal support, it too was shorter
301 than at our random sites. In this area, where the sagebrush steppe transitions to mixed
302 grass prairie, sagebrush occurs in higher densities on range sites in low seral condition and
303 soils with low productivity. Vegetative productivity on these sites can be reduced by over
304 grazing by livestock (Natural Resources Conservation Service, Electronic Field Office
305 Technical Guide, http://efotg.nrcs.usda.gov/efotg_locator.aspx?map=ND, accessed 25
306 November 2008). Thus, it is not surprising that taller grass and taller shrubs (sagebrush)
307 occurred on sites where sagebrush was less prominent. Despite the nuances of site
308 characteristics, sage-grouse appeared to select for greater concealment at and near the nest
309 and other than vegetative cover nest area variables were not important to nest selection by
310 sage-grouse in our area.

311 *Nest survival*

312 Across the range, sage-grouse nest survival averages just under 50% in relatively
313 unaltered habitats and below 40% in altered habitats (Connelly et al. 2004). Nest success
314 in stable populations generally tends to be higher (Aldridge & Brigham 2001). The
315 population of sage-grouse we studied would likely be considered unstable (declining) from
316 1951 to present (unpublished spring sage-grouse census data, North Dakota Game and Fish
317 Department, Bismarck, ND). Most of the area has been altered by historical grazing, and
318 oil and gas development. Nest survival in our study was typical of other altered habitats
319 and most nests were lost to predation. While marking nests and repeat visitations to nests
320 could attract predators, we visited nest sites only once to estimate the stage of incubation

321 and otherwise stayed >20 m away and do not believe that inconspicuous flagging on
322 sagebrush 20 m from nests increased nest predation. Nest predation can be higher in
323 fragmented landscapes (Herkert 1994, Sievert & Lloyd 1985). In the Powder River Basin
324 of Wyoming, extensive large-scale modifications of sagebrush habitat, and range
325 modification for livestock from oil and gas development were associated with significant
326 reductions in sage-grouse populations (Walker et al. 2007).

327 Of the vegetative characteristics identified in the resource selection models for
328 nests, only nest VOR was included in models of nest survival; and it had a negative
329 relation to daily survival rates. The height of the shrubs that nests were located beneath
330 was also lower at nests that survived than those that failed. We attribute these counter
331 intuitive relations to selection by sage-grouse for area of higher shrub/sagebrush which had
332 a positive influence on survival of nests. Thus, our data suggested that shrub (sagebrush)
333 cover may be more important than taller shrubs in nest survival. Areas with high shrub
334 cover also had relative low stature sagebrush. The low stature of sagebrush in these stands
335 likely resulted from the dense clay-pan soils or past grazing practices. Connelly et al. 2000
336 reviewed several studies that showed sage-grouse selecting stands with greater sagebrush
337 cover, but also the tallest sagebrush in the stand. Tall grass improved daily survival rates
338 and seemingly compensated to some degree for the shorter concealment by shrubs. When
339 shrub cover was less than about 9 - 10% or grass height was less than about 16 - 18 cm
340 nest survival declined rapidly. Although shrub cover was lower than most other studies,
341 the grass height at nest sites in our study was reflective of other studies.

342 Predators with a keen sense of smell use olfactory cues to locate nests (Storaas
343 1988), and birds that are wet have been hypothesized to have stronger odor because water
344 on the skin activates bacteria (Syrotuck 2000). Daily precipitation events decreased daily
345 survival rates which declined in a near linear manner with increasing amounts of
346 precipitation. Precipitation during incubation increased predation of wild turkey nests
347 (Roberts et al. 1995, Roberts & Porter 1998, Lehman et al. 2008). However, the relation
348 between precipitation and nest survival may be complicated by high nest attendance by the
349 female and decreased predator activity during precipitation followed by a lag effect on
350 subsequent days when females are away from the nest (Moynahan et al. 2007). However,
351 the lag of precipitation was not important in our study.

352 **Management implications**

353 If the BLM could increase its management buffer from the current 3.2 km (Bureau
354 of Land Management, 1988, Resource Management Plan, Dickenson, North Dakota) to 5
355 km, this larger area would encompass 86% of nests. Currently, there are no management
356 regulations that pertain to sage-grouse on state owned land in North Dakota. We believe a
357 strategy similar to the BLM would be beneficial to sage-grouse. Our models suggest that
358 patches of shrubs/sagebrush with > 9% canopy cover and grass taller than 16 cm improved
359 the chances of a sage-grouse surviving to hatch. Sagebrush patches selected by nesting
360 females were grazed and showed evidence of being in low seral condition due to past
361 grazing or soil characteristics. There is strong evidence in the literature that if areas of
362 sagebrush cover (>9%) occurred with taller shrubs, sage-grouse would use them and it
363 should improve nest survival. Tall grass is also important to sage-grouse nest survival and

364 grazing management that provides grass heights >16 cm should benefit nest survival. Our
365 results emphasize the importance of considering local conditions in the management of
366 sage-grouse.

367 **Acknowledgments**

368 Funding for this study was provided by Federal Aid in Wildlife Restoration Act
369 (W-67-R) through North Dakota Game and Fish Department, Bureau of Land Management
370 (ESA000013), U.S. Forest Service, Rocky Mountain Research Station
371 (05-JV-11221609-127), U.S. Forest Service Dakota Prairie National Grasslands
372 (05-CS-11011800-022), and support from South Dakota State University. Field assistance
373 was provided by A. Geigle, D. Gardner, C. Berdan, and T. Zachmeier. A number of
374 volunteers assisted during capture and radio collaring of hens and chicks. T. Apa assisted
375 with training on trapping techniques. We also acknowledge and appreciate those
376 landowners who granted us permission to conduct this study on their lands. R. King
377 provided statistical advice and support, and J. Connelly and C. Hagen provided comments
378 to earlier drafts of this manuscript.

379 **References**

- 380 Aldridge, C.L., & Brigham, R.M. 2001: Nesting and reproductive activities of Greater
381 Sage-Grouse in a declining northern fringe population. - *Condor* 103: 537-543.
- 382 Apa, A.D. 1998: Habitat use and movements of sympatric sage and Columbian sharp-
383 tailed grouse in southeastern Idaho. - PhD thesis, University of Idaho, Moscow,
384 199pp.
- 385 Barnett, J.F., & Crawford, J.A. 1994: Pre-laying nutrition of sage grouse. *Journal of*
386 *Range Management*. - 47: 114-118.
- 387 Benkobi, L., Uresk, D.W., Schenbeck, G., & King, R.M. 2000: Protocol for monitoring
388 standing crop in grasslands using visual obstruction. *Journal of Range*
389 *Management*. - 53: 627-633.
- 390 Booth, D.T., Cox, S.E., Meilke, T.W., & Fitzgerald, C. 2006: The accuracy of ground-
391 cover measurements. *Rangeland Ecology and Management*. - 59: 179-188.
- 392 Bradbury, J.W., Gibson, R.M., & McCarthy, C.E. 1989: Dispersion of displaying Sage
393 Grouse males: II, the role of female dispersion. *Behavioral Ecology of*
394 *Sociobiology*. - 24: 15-24.
- 395 Braun, C.E., Britt, T.E., & Wallestad, R.O. 1977: Guidelines for maintenance of sage
396 grouse habitats. *Wildlife Society Bulletin*. - 5: 99-106.
- 397 Burnham, K.P., & Anderson, D.R. 2002: Model selection and multimodel inference: a
398 practical information-theoretic approach, second edition. - Springer-Verlag, New
399 York, New York, 488pp.

400 Connelly, J.W., Wakkinen, W.L., Apa, A.D., & Reese, K.P. 1991: Sage grouse use of nest
401 sites in southeastern Idaho. *Journal of Wildlife Management*. - 55: 521-524.

402 Connelly, J.W., Schroeder, M.A., Sands, A.R., & Braun, C.E. 2000: Guidelines to manage
403 sage grouse populations and their habitats. *Wildlife Society Bulletin*. - 28: 967-
404 985.

405 Connelly, J.W., Knick, S.T., Schroeder, M.A., & Stiver, S.J. 2004: Conservation
406 Assessment of Greater Sage-grouse and Sagebrush Habitats. - Western Association
407 of Fish and Wildlife Agencies, Cheyenne, Wyoming, WY.

408 Cottam, G., & Curtis, J.T. 1956: The use of distance measures in phytosociological
409 sampling. *Ecology*. - 37: 451-460.

410 Crunden, C.W. 1963: Age and sex of sage grouse from wings. *Journal of Wildlife
411 Management*. - 27: 846-849.

412 Daubenmire, R.F. 1959: A canopy-coverage method of vegetation analysis. *Northwest
413 Science*. - 33: 224-227.

414 Dinsmore, S. J., White, G.C., & Knopf, F.L. 2002: - Advanced techniques for modeling
415 avian nest survival. *Ecology*. - 83: 3476-3488.

416 Floyd, D.A., & Anderson, J.E. 1987: A comparison of three methods for estimating plant
417 cover. *Journal of Ecology*. - 75: 221-228.

418 Gregg, M.A., Crawford, J.A., Drut, M.S., & DeLong, A.K. 1994: Vegetational cover and
419 predation of sage grouse nests in Oregon. *Journal of Wildlife Management*. - 58:
420 162-166.

421 Gregg, M.A., Dunbar, M.R., Crawford, J.A., & Pope, M.D. 2006: Total plasma protein
422 and reneating by greater sage-grouse. *Journal of Wildlife Management*. - 70:
423 472-478.

424 Guthery, F.S., Brennan, L.A., Peterson, M.J., & Lusk, J.J. 2005: Information theory in
425 wildlife science: critique and viewpoint. *Journal of Wildlife Management*. - 69:
426 457-465.

427 Hagen, C.A., Connelly, J.W., & Schroeder, M.A. 2007: A meta-analysis of greater
428 sage-grouse *Centrocercus urophasianus* nesting and brood-rearing habitats.
429 *Wildlife Biology*. - 13: 42-50.

430 Hays, H., Lecroy, M. 1971: Field criteria for determining incubation stage in eggs of the
431 common tern. *Wilson Bulletin*. - 83: 425-429.

432 Herkert, J.R. 1994: The effects of habitat fragmentation on Midwestern grassland bird
433 communities. *Ecological Applications*. - 4: 461-471.

434 Hoffman, R.W., Luttrell, M.P., & Davidson, W.R. 1996: Reproductive performance of
435 Merriam's wild turkeys with suspected *Mycoplasma* infection. *Proceedings of the*
436 *National Wild Turkey Symposium*. - 7: 145-151.

437 Holloran, M.J., & Anderson, S.H. 2005: Spatial distribution of greater sage-grouse nests
438 in relatively contiguous sagebrush habitats. *Condor*. - 107: 742-752.

439 Hosmer, D.W., & Lemeshow, S. 2000: *Applied Logistic Regression*, 2nd Edition. - John
440 Wiley & Sons Incorporated Publication, New York, New York, 373pp.

441 Johnson, J.R., & Larson, G.E. 1999: Grassland plants of South Dakota and the northern
442 Great Plains. - South Dakota Agricultural Experiment Station B566 (rev), South
443 Dakota State University, Brookings, 288.

444 Klebenow, D.A. 1969: Sage grouse nesting and brood habitat in Idaho. *Journal of*
445 *Wildlife Management.* - 33: 649-662.

446 Lehman, C.P., Rumble, M.A., Flake, L.D., & Thompson, D.J. 2008: Merriam's turkey
447 nest survival and factors affecting nest predation by mammals. *Journal of Wildlife*
448 *Management.* - 72: 1765-1774.

449 McCarthy, J.J., & Kobriger, J.D. 2005: Management plan and conservation strategies for
450 greater sage-grouse in North Dakota. - North Dakota Game and Fish Department,
451 Bismarck, ND, ivpp..

452 Mielke, P.W., Jr., & Berry, K.J. 2001: *Permutation Methods: a distance function*
453 *approach.* - Springer-Verlag, New York, NY, 352pp.

454 Moynahan, B.J., Lindberg, M.S., Rotella, J.J., & Thomas, J.W. 2007: Factors affecting
455 nest survival of greater sage-grouse in northcentral Montana. *Journal of Wildlife*
456 *Management.* - 71: 1773-1783.

457 North Dakota Agricultural Weather Network. 2006. *Climatology Information*, Bowman
458 ND. <http://ndawn.ndsu.nodak.edu>. (Accessed 5 Sept 2006).

459 Oyler-McCance, S.J, Taylor, S.E., & Quinn, T.W.. 2005: A multilocus population genetic
460 survey of the greater sage-grouse across their range. *Molecular Ecology.* - 14:
461 1293-1310.

462 Patterson, R.L. 1952: The sage grouse of Wyoming. - Sage Books, Inc., Denver, CO,
463 341pp.

464 Petersen, B.E. 1980: Breeding and nesting ecology of female Sage Grouse in North Park,
465 Colorado. - MS thesis, Colorado State University, Fort Collins, CO, 86pp.

466 Pollard, J.H. 1971: On distance estimators of density in randomly distributed forests.
467 Biometrics. - 27: 991-1002.

468 Robel R.J., Briggs, J.N., Dayton, A.D., & Hulbert, L.C. 1970: Relationships between
469 visual obstruction measurements and weight of grassland vegetation. Journal of
470 Range Management. - 23: 295-297.

471 Roberts, S.D., & Porter, W.F. 1998: Relation between weather and survival of wild turkey
472 nests. Journal of Wildlife Management. - 62: 1492-1498.

473 Roberts, S.D., Coffey, J.M., & Porter, W.F. 1995: Survival and reproduction of female
474 wild turkeys in New York. Journal of Wildlife Management. - 59: 437-447.

475 SAS Institute. 2005: SAS: JMP Start Statistics. – SAS Publishing Incorporated, New
476 York, NY.

477 Schroeder, M.A. 1997: Unusually high reproductive effort by Sage Grouse in a
478 fragmented habitat in north-central Washington. Condor. - 99: 933-941.

479 Schroeder, M.A., Aldridge, C.L., Apa, A.D., Bohne, J.R., Braun, C.E., Bunnell, S.D.,
480 Connelly, J.W., Diebert, P.A., Gardner, S.C., Hilliard, M.A., Kobriger, G.D., &
481 McCarthy, C.W. 2004: Distribution of Sage Grouse in North America. Condor. -
482 106: 363-376.

483 Sievert, P. R., & Lloyd, K.B. 1985: Survival of Snowshoe Hares at a Geographical
484 Boundary. *Journal of Wildlife Management*. - 49: 854-866.

485 Smith, J.T., Flake, L.D., Higgins, K.F., Kobriger, G.D., & Homer, C.G. 2005: Evaluating
486 lek occupancy of greater sage-grouse in relation to landscape cultivation in the
487 Dakotas. *Western North American Naturalist*. - 65: 310-320.

488 SPSS. 2002: SPSS for Windows 11.5. - SPSS, Inc. Chicago, IL.

489 Stephens, P.A., Buskirk, S.W., Hayward, G.D., & del Rio, C.M.. 2005: Information theory
490 and hypothesis testing: a call for pluralism. *Journal of Applied Ecology*. - 42: 4-12.

491 Storaas, T. 1988: A comparison of losses in artificial and naturally occurring capercaillie
492 nests. *Journal of Wildlife Management*. - 52: 123-126.

493 Sveum, C.M. 1995: Habitat selection by Sage Grouse hens during the breeding season in
494 south-central eastern Washington. - MS thesis, Oregon State University, Corvallis,
495 OR, 86pp.

496 Sveum, C.M., Edge, W.D., & Crawford, J.A. 1998: Nesting habitat selection by sage
497 grouse in southcentral Washington. *Journal of Range Management*. - 51: 265-269.

498 Syrotuck, W.G. 2000: Scent and the scenting dog. - Barkleigh Productions, Inc.,
499 Mechanicsburg, PA

500 Wakkinen, W.L. 1990: Nest site characteristics and spring-summer movements of
501 migratory Sage Grouse in southeastern Idaho. - M.S. thesis, University of Idaho,
502 Moscow, ID, 57pp.

503 Wakkinen, W.L., Reese, K.P., & Connelly, J.W. 1992a: Sage grouse nest location in
504 relation to leks. *Journal of Wildlife Management*. - 56: 381-383.

- 505 Wakkinen, W.L., Reese, K.P., Connelly, J.W., & Fischer, R.A. 1992b: An improved
506 spotlighting technique for capturing Sage Grouse. *Wildlife Society Bulletin*. - 20:
507 425-426.
- 508 Walker, B.L., Naugle, D.E., & Doherty, K.E. 2007: Greater sage-grouse population
509 response to energy development and habitat loss. *Journal of Wildlife Management*.
510 - 71: 2644-2654.
- 511 Wallestad, R.O., & Pyrah, D.B. 1974: Movement and nesting of sage grouse hens in
512 central Montana. *Journal of Wildlife Management*. - 38: 630-633.
- 513 Welch, B.L. 2005: Big sagebrush: as sea fragmented into lakes, ponds, and puddles. - U.S.
514 Forest Service, Rocky Mountain Research Station, General Technical Report
515 RMRS-GTR-144, Fort Collins, CO, 210pp.
- 516 White, G.C., & Burnham, K.P. 1999: Program MARK: survival estimation from
517 populations of marked animals. *Bird Study*. - 46: 120-138.

518 Table 1. Average of key vegetative (\pm SD) characteristics from greater-sage grouse nest
 519 sites and random sites included in resource selection models from North Dakota,
 520 USA, 2005-2006.

Vegetative Characteristic	Nest		Random		P-value ¹
	\bar{x}	\pm SD	\bar{x}	\pm SD	
Percent total vegetative cover	70.4	15.5	54.4	20.4	>0.01
Percent total grass cover	27.4	13.6	19.3	14.6	0.01
Percent total forb cover	15.4	11.8	11.0	6.6	0.05
Percent shrub cover ²	9.8	4.0	7.1	4.6	>0.01
Percent litter	12.9	8.3	7.9	5.0	0.01
Percent intermediate wheatgrass	1.7	2.3	2.4	2.1	0.06
Percent green needlegrass	2.5	3.3	1.3	1.8	0.06
Percent western wheatgrass	4.2	3.7	2.1	2.6	<0.01
Percent Kentucky bluegrass	3.3	3.7	1.9	2.7	0.05
Nest VOR (inches)	2.6	1.2	2.1	1.6	0.02
1-m VOR	3.9	2.1	2.7	2.3	>0.01
2-m VOR	3.0	2.1	2.73	2.3	<0.01
3-m VOR	2.5	1.4	2.2	1.8	0.12
4-mVOR	2.4	1.5	2.1	1.8	0.30
5-m VOR	2.2	1.3	2.2	2.1	0.24
Site VOR	2.6	1.2	2.1	1.6	0.02
Sagebrush density (ha)	2,576.1	1,833.6	1,399.4	1,795.1	>0.01

521

522 Table 1. continued.

523

Grass height (inches)	10.2	3.7	11.2	3.3	0.17
Shrub height (mm)	42.1	18.4	48.4	16.7	0.07

524

525 ¹ From multiple response permutation procedure (Mielke and Berry 2001).

526 Table 2. Summary of model selection of logistic regression for greater sage-grouse nests ($n = 34$) from random sites ($n = 50$) in
 527 North Dakota, USA, 2005-2006 using the Information Theoretic approach.

Model ¹	AIC	ΔAIC_c	$AICw_i$	K^2	Deviance
Total cover + shrub height + nest VOR	95.14	0.00	0.23	5	84.37
Total cover + shrub height + 1-m VOR	95.52	0.39	0.18	5	84.76
Total cover + shrub height	96.17	1.03	0.13	4	87.66
Total cover + shrub height + 1-m VOR + grass height	97.113	1.97	0.08	6	84.02
Total cover + shrub height + nest VOR + grass height	97.12	1.98	0.08	6	84.03
Global model	104.48	9.34	<0.01	10	81.47

528
 529 ¹ A total of 25 models were considered. Model results are presented in descending order of rankings and include models with
 530 ΔAIC values less than 2.0.

531 ² Number of parameters includes those in model plus year, and intercept.

532 Table 3. Summary of model selection for greater sage-grouse nest survival considering vegetation characteristics and time-
 533 dependent variables in southwestern North Dakota, USA, 2005-2006.

Models ¹	AICc ²	Δ AICc	AICw ³ _i	K ⁴	Deviance
Grass height + shrub cover + nest VOR + precipitation	109.537	0	0.291 (0.376)	5	99.38
Grass height + shrub cover + total cover + nest VOR + precipitation	110.038	0.50	0.226	6	97.82
Grass height + shrub cover + nest VOR + site VOR + precipitation	111.129	1.59	0.13 (0.170)	6	105.45
Grass height + shrub cover + nest VOR	113.565	4.02	0.039	4	105.46
Precipitation	115.888	6.35	0.014	2	111.86
Constant survival	119.025	9.49	0.002	1	117.02
Nest age	120.219	10.68	0.001	2	116.19
Year	120.894	11.38	0.001	2	116.86
Best nest site selection model: total cover + shrub height + nest VOR	122.586	13.319	<0.001	4	114.751

534 ¹ A total of 41 models were evaluated. We evaluated vegetation characteristics first, then included precipitation and
 535 temperature. The first 3 models were the highest ranked models with or without precipitation.

536 ² Methods and interpretation of heading are described by Burnham and Anderson (2002).

537 ³ AIC_w in parentheses are those with the second model eliminated.

538 ⁴ Number of parameters in the model includes the intercept from constant survival estimate.

539 Figure 1. Daily survival as a function of A) precipitation and percent shrub cover with nest VOR and grass height constant at
540 mean, B) precipitation and grass height with nest VOR and percent shrub cover held constant at the mean, and C) Grass height
541 and shrub cover with precipitation and nest VOR held constant and the mean.

542

