

1 **Hunting bag and distance from nearest day-roost in Camargue ducks**
2

3 Matthieu Guillemain ^a, Jean-Yves Mondain-Monval ^a, Emilien Weissenbacher ^a, Anne-Laure
4 Brochet ^{a,b} & Anthony Olivier ^b
5

6 ^a *Office National de la Chasse et de la Faune Sauvage, CNERA Avifaune Migratrice, La Tour*
7 *du Valat, Le Sambuc, 13200 Arles, France.*

8 ^b *La Tour du Valat, Le Sambuc, 13200 Arles, France.*
9

10

11

12

13 ***Running head:*** Guillemain *et al.*: Duck hunting bag in Camargue.

14

15 ***Keywords:*** Hunting bag, day-roost, Camargue, winter, distance, wildfowl.
16

17

18

19 Corresponding Author :

20 Matthieu Guillemain

21 *Office National de la Chasse et de la Faune Sauvage, CNERA Avifaune migratrice, La Tour*
22 *du Valat, Le Sambuc, 13200 Arles, France.*

23 Tel : +33. 490 97 27 87.

24 Fax: +33 490 97 27 88.

25 e-mail : matthieu.guillemain@oncfs.gouv.fr
26
27
28

1 **Abstract**

2

3 Reserves that are closed to hunting are important to wintering wildfowl (*Anatidae*). Reserves
4 can also benefit local hunters because they are often used as day-roosts by birds prior to
5 nocturnal feeding in surrounding wetlands. In this study, we used annual hunting bags from
6 45 hunting estates in the Camargue, southern France, to study the relationship between
7 hunting success and distance to the nearest duck day-roost, which are generally located in
8 protected areas. Five dabbling duck species (*Anas* spp.) and a diving duck species (pochard
9 *Aythya ferina*) were studied. The relationship between hunting bag and distance from roost
10 was negative in all cases except mallard (*Anas platyrhynchos*), which is subject to specific
11 hunting management practices (i.e. raise and release). A significance threshold was not
12 reached in wigeon (*A. penelope*) and shoveler (*A. clypeata*). This supports the hypothesis that
13 hunting estates closer to protected areas are more successful; however, the decrease in hunting
14 bags with increasing distance from the nearest roost could be seen in as little as a few
15 thousands of meter in all species. This suggests that only the hunting grounds with a distance
16 less than 2-3 kilometres from the nearest roost may actually have increased hunting
17 accessibility for certain duck species.

1 **Introduction**

2 Wildfowl (*Anatidae*) are sensitive to human activities during the hunting season (e.g. Bell &
3 Owen 1990, Tamisier et al. 2003, Triplet et al. 2003, Blanc et al. 2006) and often gather in
4 large numbers in disturbance-free protected areas (Cox & Afton 1997, Fox & Madsen 1997,
5 Madsen, Pihl & Clausen 1998). Refuges can also benefit hunters when they are used by birds
6 as roosts. Ducks departing from refuges for nocturnal feeding or other activities are more
7 exposed to hunting activities (Bellrose 1954, Griffith 1957, Anonymous 1961, Bell & Owen
8 1990, Mathevet & Tamisier 2002). Some duck species commute two times within a 24-hour
9 period, moving from day-roosts to nocturnal foraging areas. These two habitats are considered
10 as their functional unit (Tamisier 1978). Regular commuting patterns during dusk and dawn
11 increase the ducks vulnerability to hunting. Different studies have shown that the average
12 distance ducks fly between roosting and foraging areas varies among species and
13 environmental conditions (e.g. adverse weather) and can range from 0.8 to 50 km (Fog 1958,
14 Frazer et al. 1990). However, it seems intuitive that ducks would minimize travel distance
15 given the energetic cost of flight. Distance between day-roosts and nocturnal foraging habitats
16 is therefore generally limited to a few kilometres (Tamisier & Tamisier 1981, Jorde et al.
17 1983, Guillemain et al. 2002, Legagneux et al. unpubl. data). It is hypothesised that hunting
18 areas adjacent to day-roosts, whether protected or not, experience greater hunting harvests.
19 One study in Camargue in the south of France demonstrated that habitats protected from
20 hunting and comprising a major roost influenced surrounding land management practices,
21 which became more hunting-orientated. As a result, the price of hunting leases in the vicinity
22 of the reserve increased to more than 1800 Euros per hunter per year (Mathevet & Tamisier
23 2002). More information concerning duck movements is necessary to assess the distance
24 travelled in regards to land management procedures. Knowledge of duck movements could
25 influence decisions to protect or manage nocturnal foraging habitats in addition to day-roosts,

1 or to establish buffer areas around reserves (Fox & Madsen 1997, Rodgers & Smith 1997).
2 This information could also be important for understanding the local dispersal of propagules
3 (seeds or invertebrates [Green et al. 2002]) or the spread of diseases by ducks, such as avian
4 influenza viruses (Gauthier-Clerc et al. 2007, Saad et al. 2007 for teal).

5 Our research evaluated hunters' harvests from various private hunting estates in the
6 Camargue, southern France, to test the relationship between hunters' harvests and distance
7 from hunting estates and nearest day-roosts.

8

9 **Study area and methods**

10 The Camargue encompasses 145,000ha of the Rhône Delta (see Tamisier & Dehorter 1999
11 for a general description). The Camargue is a site of international importance for waterbirds
12 and exceeds the criterion of international importance for several duck species (Scott & Rose
13 1996). More than 85,000ha of the delta are wetlands, and 78% of the wetlands are divided
14 into approximately 180 private estates that integrate hunting activities (Dehorter & Tamisier
15 1996). As part of a research program on hunting practices and wildfowl population dynamics
16 in the Camargue (Mondain-Monval et al. 2006), hunting bags are collected annually for as
17 many estates as possible. In the present study, we used data from 45 estates which provided
18 duck bags for at least 3 hunting seasons, together with an indication of hunting pressure
19 (number of hunter-days). Data ranged from the 1926-1927 to the 2004-2005 hunting seasons.
20 However, most of the estates provided datasets beginning in the 1980's or 1990's (Appendix
21 1), which allows for comparison by site and also relates to the period when the list of duck
22 day-roosts were established by aerial counts (see below). The number of annual data ranged
23 from 3 to 68 years between the 45 estates (median = 9 years). The inclusion of the 13 estates
24 with <5 years of data led to a 22 to 33% increase in the variance of the daily bag per hunter,
25 depending on duck species, compared to the reduced dataset made of the 32 estates with 5

1 years of data or more. However, the results concerning the relationship between daily bag and
2 distance from the nearest roost were exactly the same in terms of significant and non-
3 significant patterns. The complete 45 estate dataset was thus used to cover the largest
4 geographical area possible. Datasets were interrupted in some cases, due to changes in
5 landlords and/or hunting lease holders. Three datasets also abruptly ended when the hunting
6 estates became natural reserves. In each estate, the average daily bag per duck species per
7 hunter was calculated for each year, and then averaged for all available years (Appendix 1).

8 Although owners of hunting estates sometimes manage water levels, salinity and frequency
9 of disturbance so as to promote duck roosting within their property, most ducks spent daylight
10 hours in the over 20,000 ha of protected areas in the Camargue. Based on long-term (1964-65
11 to 1994-95) aerial monitoring of wintering wildfowl, Tamisier & Dehorter (1999, pp. 352-
12 353) provided a list of the main day-roosts for the different duck species based on the average
13 number of birds of each species they hosted over the monitoring period. We determined the
14 central point of each of these roosts for each species through GIS software (Arcview 3.1 GIS,
15 ESRI 1998). The central point of each of the 45 hunting estates was also computed using the
16 same methodology, allowing calculation of the distance in kilometres between each estate and
17 the nearest day-roost for each duck species (NB: the nearest day-roost was possibly different
18 for different duck species in a given hunting estate). This method therefore established a
19 distance equal to zero in the case of day-roosts within hunting estates.

20 The distribution of hunting bag values was non-normal (Kolmogorov-Smirnov tests: all d
21 values > 0.20 , all $P < 0.01$), and usual transformations (such as logarithm, square-root, etc)
22 did not solve this problem. Given the distribution of the data, we used maximum-likelihood
23 ratio statistics based on quasi-Poisson distribution of the variables. The relationship between
24 hunting bag and distance from the nearest roost in each duck species was analysed with a non-
25 linear regression model, using a negative exponential relationship. Pintail (*Anas acuta*), red-

1 crested pochard (*Netta rufina*) and tufted duck (*Aythya fuligula*) were not included in the
2 analyses because they were either absent or represented only a small part of the total bag of
3 the 45 studied hunting estates (0.005 to 0.063 individuals per hunter per day on average on 20
4 estates, and 25 estates with no single individual ever harvested in Tufted Duck). The dataset
5 therefore included mallard (*Anas platyrhynchos*), teal, gadwall (*Anas strepera*), shoveler
6 (*Anas clypeata*), wigeon (*Anas penelope*) and pochard (*Aythya ferina*) hunting bags.

7

8 **Results**

9 All six duck species except mallard showed the same pattern of decreasing daily bag per
10 hunter with increasing distance from the nearest day-roost, although this was non-significant
11 in shoveler and marginally so in wigeon (Fig. 1, Table 1). All species' hunting bags were
12 highly variable among estates close to day-roosts. The range was as wide as 0.1 to 4.5 teal per
13 hunter per day for estates less than 2 km from a day-roost. The lack of significance in
14 shoveler and wigeon was essentially due to the high variance in hunting bags for estates close
15 to day-roosts (0 to 0.8 Wigeon per hunter per day and 0 to 1.3 Shoveler per hunter per day for
16 estates < 1 km from nearest roost). However, the wigeon and shoveler average daily hunting
17 bag also tended to decrease exponentially with increasing distance from the roost (Table 1).

18 In all cases the decrease in hunting bag with increasing distance was very rapid, since
19 estates further than 5 km from a roost harvested very limited numbers of individuals of a
20 given species (Fig. 1). Using the significant trends for the teal, gadwall and pochard models
21 (see Table 1), we calculated the average expected daily bag per kilometre of distance from the
22 nearest roost and determined the threshold distance above which this value no longer
23 exceeded 10% of the expected bag within the first kilometre. In the three species, the
24 threshold values were 9, 5 and 3 kilometres, respectively (i.e. an estate located more than 3

1 km from any pochard day-roost could not expect a daily bag per hunter higher than 10% of
2 the daily bag of an estate within 1 km from such a roost).

3

4 **Discussion**

5 All daily hunting bags decreased (though a significant threshold was not reached in Wigeon
6 and Shoveler) in an exponential manner with increasing distance from the nearest day-roost
7 with the exception of mallard. This is consistent with central place foraging and refuge
8 theories, in which animals try to limit their travelling distance between a central protected
9 area and their feeding areas (e.g. review in Stephens & Krebs 1986, Cox & Afton 1996). The
10 fact that daily bags differed among estates less than one kilometre from a roost probably
11 reflects management options as well as the differences in hunting estate sizes. Estates are
12 managed and decisions are made depending on the priority species that the managers aim to
13 attract (e.g. by selecting different water regimes, water levels or salinity) and the habitat
14 constraints (e.g. water salinity). For example, if an estate is close to both a teal and a pochard
15 day-roost, the management options and habitat will favour either one or the other species.
16 Furthermore, some hunting estates adjacent to or encompassing day-roosts allow shooting
17 during a few daylight hours during the morning (as opposed to dusk and dawn shooting
18 during duck commuting flights between day-roost and nocturnal feeding areas), which is
19 obviously a more efficient hunting technique. Depending on species, ducks virtually
20 disappeared from hunting bags in estates located more than 3-10 km from the day-roosts and
21 in all cases were mostly shot within 2-3 km range. It is interesting to note that these distances
22 between roosts and selected foraging areas are within the same ranges determined from earlier
23 studies using radio-tracking methodologies. Radio-tracking studies suggest that ducks
24 generally travel from a few hundred meters up to 10 km to feed, this range being species-
25 specific (Tamisier & Tamisier 1981, Jorde et al. 1983, Frazer et al. 1990, Cox & Afton 1996,

1 Guillemain et al. 2002, Legagneux et al. unpubl. data). The present study suggests that larger
2 home ranges (including up to 50 km travels) (e.g. Bellrose 1954, Fog 1968, references in
3 Jorde et al. 1983) probably represented extreme travel distances or areas where no suitable
4 habitat was available to the birds within their preferred travel distance. Some of these
5 previous studies considered ring recoveries, indicating that the distance travelled may have
6 been over several days.

7 The insignificant relationship between daily bag and distance from roost in mallard was
8 unexpected, but could be explained by hunting management practices for this species. Most
9 mallards in the Camargue are harvested in the beginning of the season and many of which are
10 young individuals hatched on the estate during the previous spring or raised and released in
11 spring and summer, a common practice in France. The exact number of such released mallard
12 is not known but has been estimated to be around 20-30,000 individuals in the Camargue
13 (Tamisier 2004). The uptake of locally-released or locally-hatched individuals within the
14 hunting estates at the beginning of the season may therefore hide a potential relationship
15 between hunting bag and distance from a roost that may occur in natural wild conditions. This
16 practice is, however, unlikely to have affected the uptake of other ducks, since the average
17 daily bag in estates releasing ($n = 10$ properties) or not releasing mallard ($n = 35$ properties)
18 did not differ significantly in any species (t tests, all t absolute values < 1.33 , all $P > 0.1917$).

19 From a hunting management point of view, this study suggests that the management
20 practices leading to roost establishments within the hunting estate may indeed lead to larger
21 daily bags (especially if this allows diurnal shooting). Similarly, the present results show that
22 the higher prices of hunting leases around protected duck day-roosts observed by Mathevet &
23 Tamisier (2002) can be explained by higher expected shooting opportunities in adjacent
24 estates. However, private properties are not affected by roosts located more than 2-3 km
25 away. On the other hand, we suggest that where the protection of duck species is a priority

1 over hunting considerations, the establishment of hunting-free buffer areas around core
2 wildfowl refuges (e.g. Fox & Madsen 1997) could be an effective strategy since the radius of
3 such buffer zones may be as little as a few kilometres (as opposed to 20-50 kilometres in
4 some earlier studies) for most species.

5 If they do disperse seeds, invertebrate eggs or disease locally during their feeding travels,
6 the management implications of our results are that dabbling and diving ducks are most likely
7 to do so over short distances during their daily movements. This does not take into
8 consideration the long distances they may cover during migration which can be associated
9 with long-distance dispersal of seeds or invertebrates (Green & Figuerola 2005). This
10 conclusion is only based on indirect information provided by birds shot by hunters, therefore
11 future research using electronic devices to determine actual nocturnal movements of ducks
12 should be carried out. This may also allow for future testing to determine if wintering
13 wildfowl actually have longer daily travel distances in order to avoid hunted estates and move
14 between protected areas as observed in the past in western France (Guillemain et al. 2002).

15
16
17

18 **Acknowledgements**

19 We are grateful to the many Camargue landowners and holders of hunting leases who
20 provided information on their annual hunting bags. We would also like to thank Michel
21 Gauthier-Clerc, Andy Green, Jesper Madsen, Jean-Baptiste Mouronval and Lisa Ernoul for
22 their useful comments on an earlier version of the manuscript, as well as Hervé Fritz for
23 valuable statistical advice. Anne-Laure Brochet is funded by a Doctoral grant from Office
24 National de la Chasse et de la Faune Sauvage, and receives additional funding from a research
25 agreement between ONCFS, the Tour du Valat and the Laboratoire de Biométrie et de
26 Biologie Evolutive UMR 5558 CNRS Université Lyon 1 in France, together with the Doñana
27 Biological Station (CSIC) in Spain. This work also received funding from the Agence

- 1 Nationale de la Recherche through the Santé Environnement - Santé Travail scheme (contract
- 2 number 2006-SEST-22).
- 3

1 **References**

- 2 Anonymous 1961: National wildfowl refuges. - *Wildfowl* 12: 23-25.
- 3 Bell, D.V. & Owen, M. 1990: Shooting disturbance – a review. – In: Matthews, G.V.T. (Ed.);
4 Managing waterfowl populations. IWRB Spec. Pub. 12. IWRB, Slimbridge, pp. 159-171.
- 5 Bellrose, F.C. 1954: The value of waterfowl refuges in Illinois. - *Journal of Wildlife*
6 *Management* 18: 160-169.
- 7 Blanc, R., Guillemain, M., Mouronval, J.B., Desmots, D. & Fritz, H. 2006: Effects of non-
8 consumptive leisure disturbance to wildlife. – *Revue d'Ecologie (Terre & Vie)* 61: 117-
9 133.
- 10 Cox, R.R.Jr. & Afton, A.D. 1996: Evening flights of female Northern Pintails from a major
11 roost site. - *Condor* 98: 810-819.
- 12 Cox, R.R.Jr. & Afton, A.D. 1997: Use of habitats by female Northern Pintails wintering in
13 Southwestern Louisiana. - *Journal of Wildlife Management* 61: 435-443.
- 14 Dehorter, O. & Tamisier, A. 1996: Wetland habitat characteristics for waterfowl wintering in
15 Camargue, Southern France: implications for conservation. - *Revue d'Ecologie (Terre &*
16 *Vie)* 51: 161-172.
- 17 ESRI 1998. Arc View GIS. Environmental Systems Research Institute.
- 18 Fog, J. 1968: Krikandens (*Anas crecca*) spredning under fourageringstogter fra en rasteplass
19 (Albuebugten vildtreservat, Fano). (English summary). - *Dansk Ornitologisk Forenings*
20 *Tidsskrift* 62: 32-36.
- 21 Fox, A.D. & Madsen, J. 1997: Behavioural and distributional effects of hunting disturbance
22 on waterbirds in Europe: implications for refuge design. - *Journal of Applied Ecology* 34:
23 1-13.
- 24 Frazer, C., Longcore, J.R. & McAuley, D.G. 1990: Habitat use by postfledging American
25 black ducks in Maine and New Brunswick. - *Journal of Wildlife Management* 54: 451-459.

- 1 Gauthier-Clerc, M., Lebarbenchon, C. & Thomas, F. 2007: Recent expansion of highly
2 pathogenic avian influenza H5N1: a critical review. - *Ibis* 149: 202-214.
- 3 Green, A. J. & Figuerola, J. 2005: Recent advances in the study of long-distance dispersal of
4 aquatic invertebrates via birds. - *Diversity and Distributions* 11: 149-156.
- 5 Green, A. J., Figuerola, J. & Sanchez, M. I. 2002: Implications of waterbird ecology for the
6 dispersal of aquatic organisms. - *Acta Oecologica-International Journal of Ecology* 23:
7 177-189.
- 8 Griffith, R.E. 1957: Management of waterfowl habitat on the National Wildlife Refuges. -
9 *Transactions of the North American Wildlife Conference* 22: 215-220.
- 10 Guillemain, M., Fritz, H. & Duncan, P. 2002 : The importance of protected areas as nocturnal
11 feeding grounds for dabbling ducks wintering in western France. - *Biological Conservation*
12 103: 183-198.
- 13 Jorde, D.G., Krapu, G.L. & Crawford, R.D. 1983: Feeding ecology of Mallards wintering in
14 Nebraska. - *Journal of Wildlife Management* 47: 1044-1053.
- 15 Madsen, J., Pihl, S. & Clausen, P. 1998: Establishing a reserve network for waterfowl in
16 Denmark: a biological evaluation of needs and consequences. - *Biological Conservation*
17 85: 241-255.
- 18 Mathevet, R. & Tamisier, A. 2002: Creation of a nature reserve, its effect on hunting
19 management and waterfowl distribution in the Camargue (southern France). - *Biodiversity*
20 *and Conservation* 11: 509-519.
- 21 Mondain-Monval, J.-Y., Defos du Rau, P., Desnouhes, L., Mathon, N. & Olivier, A.
22 2006. The monitoring of hunting bags and hunting effort in the Camargue,
23 France. In: Boere, G.C., Galbraith, C.A. & Stroud D.A. (eds) 2006.
24 *Waterbirds around the World*. The Stationery Office, Edinburgh, UK. Pp.
25 862-864.

- 1 Rodgers, J.A. & Smith, H.T. 1997: Buffer zone distances to protect foraging and loafing
2 waterbirds from human disturbance in Florida. - *Wildlife Society Bulletin* 25: 139-145.
- 3 Saad, M.D., Ahmed, L.S., Gamal-Eldein, M.A., Fouda, M.K., Khalil, F.M., Yingst, S.L.,
4 Parker, M.A. & Montevillel, M.R. 2007: Possible avian influenza (H5N1) from migratory
5 bird, Egypt. - *Emerging Infectious Diseases* July 2007.
- 6 Scott, D.A. & Rose, P.M. 1996: Atlas of Anatidae populations in Africa and Western Eurasia.
7 Wetlands International Special Publication 41. Wetlands International, Wageningen, The
8 Netherlands.
- 9 Stephens, D.W. & Krebs, J.R. 1986: Foraging Theory. Princeton University Press, Princeton,
10 New-Jersey.
- 11 Tamisier, A. 1978: The functional units of wintering ducks: a spatial integration of their
12 comfort and feeding requirements. – *Verhandlungen der Ornithologischen Gesellschaft in*
13 *Bayern* 23: 229-238.
- 14 Tamisier, A. 2004: Camargue, quartier d’hiver de canards et foulques. Modalités de
15 fonctionnement, facteurs limitants et perspectives d’avenir. In Isenmann, P. (Ed.); *Les*
16 *oiseaux de Camargue et leurs habitats – Une histoire de cinquante ans 1954-2004*. Buchet-
17 Chastel, Paris, pp. 149-204.
- 18 Tamisier, A. & Tamisier, M.C. 1981: L’existence d’unités fonctionnelles démontrée chez les
19 sarcelles d’hiver en Camargue par la biotélémétrie. - *Revue d’Ecologie (Terre et Vie)* 35:
20 563-579.
- 21 Tamisier, A. & Dehorter, O. 1999: Camargue, canards et foulques – fonctionnement et
22 devenir d’un prestigieux quartier d’hivernage. Centre Ornithologique du Gard, Nîmes,
23 France.

1 Tamisier, A., Béchet, A., Jarry, G., Lefeuvre, J.C. & LeMaho, Y. 2003: Effets du
2 dérangement par la chasse sur les oiseaux d'eau – Revue de Littérature. – Revue
3 d'Ecologie (Terre & Vie) 58: 435-449.

4 Triplet, P., Sournia, A., Joyeux, E. & Le Dréan-Quéneec'Hdu, S. 2003: Activités humaines et
5 dérangement : l'exemple des oiseaux d'eau. – Alauda 71: 305-316.

6

7

1 **Figure Legend**







2

3

4 Figure 1. Relationship between daily bag per hunter and distance between hunting estate and
5 nearest day-roost in the six duck species. See Table 1 for statistics.

1 Guillemain et al. Fig. 1

2 |



3

4

1 **Table1.** Non-linear regressions of average daily hunting bags per hunter and the distance
 2 between hunting estates and the nearest day-roost, per duck species. The equation is given
 3 where model fit is significant.

4
 5
 6

	Likelihood Ratio	CHI ²	P-value	Model equation
Teal <i>Anas crecca</i>	-51.11	9.35	0.0022	Daily Bag = EXP(0.8062-0.0003xDistance)
Gadwall <i>A. strepera</i>	-21.63	10.11	0.0015	Daily Bag = EXP(0.1393-0.0005xDistance)
Mallard <i>A. platyrhynchos</i>	-54.67	2.71	0.1000	
Shoveler <i>A. clypeata</i>	-24.16	1.78	0.1837	
Wigeon <i>A. penelope</i>	-12.91	3.47	0.0624	
Pochard <i>Aythya ferina</i>	-11.68	6.49	0.0109	Daily Bag = EXP(-0.8824-0.0011xDistance)

Appendix 1. Hunting pressure (average annual number of hunter-days) and hunting bag (average number of pieces of game per hunter per day) for each of the 45 hunting estates studied in the Camargue, in the south of France. Values are means \pm SE. The names of the estates are not provided because of confidentiality agreements with the landowners and holders of hunting leases.

Estate	No of Years	Range	Hunting Pressure (hunter.days)	Bag (number of ducks per hunter per day)					
				Teal	Gadwall	Mallard	Shoveler	Wigeon	Pochard
				<i>Anas crecca</i>	<i>Anas strepera</i>	<i>Anas platyrhynchos</i>	<i>Anas clypeata</i>	<i>Anas penelope</i>	<i>Aythya ferina</i>
Estate 1	8	1997-2004	478.4 \pm 51.0	0.55 \pm 0.07	0.30 \pm 0.03	0.95 \pm 0.08	0.28 \pm 0.04	0.16 \pm 0.02	0.04 \pm 0.01
Estate 2	4	2001-2004	209.0 \pm 3.0	0.09 \pm 0.03	0.05 \pm 0.03	0.49 \pm 0.08	0.34 \pm 0.06	0.02 \pm 0.01	0.00 \pm 0.00
Estate 3	3	2002-2004	179.7 \pm 21.4	0.15 \pm 0.04	0.01 \pm 0.00	0.52 \pm 0.09	0.07 \pm 0.02	0.02 \pm 0.00	0
Estate 4	68	1926-2004	116.2 \pm 3.6	5.89 \pm 0.35	1.80 \pm 0.18	3.31 \pm 0.24	1.27 \pm 0.09	0.63 \pm 0.09	0.36 \pm 0.08
Estate 5	10	1979-1997	389.8 \pm 24.4	0.36 \pm 0.06	0.10 \pm 0.02	0.57 \pm 0.07	0.21 \pm 0.03	0.05 \pm 0.01	0.01 \pm 0.00
Estate 6	21	1984-2004	100.5 \pm 4.2	1.62 \pm 0.20	1.91 \pm 0.11	2.08 \pm 0.26	0.44 \pm 0.06	0.07 \pm 0.01	0.78 \pm 0.24
Estate 7	26	1979-2004	108.0 \pm 6.1	4.50 \pm 0.71	0.29 \pm 0.04	3.80 \pm 0.58	0.26 \pm 0.04	0.01 \pm 0.00	0.02 \pm 0.00
Estate 8	10	1993-2004	275.0 \pm 7.9	1.80 \pm 0.24	0.93 \pm 0.15	1.30 \pm 0.21	0.46 \pm 0.08	0.07 \pm 0.02	0.98 \pm 0.17
Estate 9	3	1984-2004	554.7 \pm 121.5	0.32 \pm 0.03	0.00 \pm 0.00	0.40 \pm 0.22	0.03 \pm 0.01	0.01 \pm 0.00	0.03 \pm 0.01
Estate 10	10	1993-2004	115.2 \pm 4.5	0.17 \pm 0.03	0.30 \pm 0.08	0.83 \pm 0.11	0.09 \pm 0.03	0.08 \pm 0.02	0.04 \pm 0.03
Estate 11	10	1990-2004	391.6 \pm 53.4	1.08 \pm 0.14	0.02 \pm 0.01	0.70 \pm 0.08	0.09 \pm 0.03	0.00 \pm 0.00	0.00 \pm 0.00
Estate 12	6	1999-2004	283.3 \pm 8.3	0.73 \pm 0.09	0.13 \pm 0.02	3.51 \pm 0.18	0.12 \pm 0.03	0.05 \pm 0.02	0.01 \pm 0.01
Estate 13	29	1973-2004	438.1 \pm 4.1	1.97 \pm 0.23	1.04 \pm 0.10	1.36 \pm 0.14	0.98 \pm 0.08	0.34 \pm 0.05	0.10 \pm 0.02
Estate 14	10	1990-2004	302.1 \pm 57.4	1.33 \pm 0.46	0.09 \pm 0.02	1.82 \pm 0.57	0.12 \pm 0.03	0.00 \pm 0.00	0
Estate 15	6	1999-2004	244.5 \pm 1.5	0.98 \pm 0.19	0.49 \pm 0.10	3.22 \pm 0.30	0.54 \pm 0.07	0.30 \pm 0.03	0.02 \pm 0.00
Estate 16	15	1981-1995	192.7 \pm 23.8	1.25 \pm 0.16	1.11 \pm 0.13	1.88 \pm 0.23	0.73 \pm 0.16	0.76 \pm 0.09	0.38 \pm 0.09
Estate 17	4	2001-2004	322.5 \pm 7.5	0.46 \pm 0.11	0.01 \pm 0.01	0.57 \pm 0.09	0.09 \pm 0.05	0.01 \pm 0.00	0
Estate 18	4	1994-2004	554.0 \pm 82.1	0.80 \pm 0.17	0.01 \pm 0.00	1.16 \pm 0.32	0.11 \pm 0.02	0.05 \pm 0.02	0.01 \pm 0.00
Estate 19	8	1997-2004	356.8 \pm 25.2	0.58 \pm 0.08	0.08 \pm 0.01	0.74 \pm 0.07	0.26 \pm 0.05	0.33 \pm 0.06	0.06 \pm 0.03
Estate 20	6	1998-2004	68.2 \pm 4.1	0.39 \pm 0.11	0.01 \pm 0.01	0.64 \pm 0.13	0	0.04 \pm 0.04	0
Estate 21	4	2001-2004	393.3 \pm 45.6	0.15 \pm 0.04	0.05 \pm 0.02	1.05 \pm 0.10	0.02 \pm 0.00	0.01 \pm 0.00	0
Estate 22	17	1988-2004	506.1 \pm 23.2	1.22 \pm 0.14	0.38 \pm 0.06	4.23 \pm 0.71	0.51 \pm 0.07	0.03 \pm 0.01	0.38 \pm 0.10

Estate 23	4	2000-2003	309.3 ± 9.9	0.21 ± 0.08	0.01 ± 0.01	0.39 ± 0.13	0.08 ± 0.04	0.00 ± 0.00	0
Estate 24	4	2001-2004	107.8 ± 18.7	0.09 ± 0.05	0.06 ± 0.02	1.22 ± 0.31	0.01 ± 0.01	0.00 ± 0.00	0
Estate 25	3	1997-2004	318.0 ± 114.0	0.11 ± 0.06	0.01 ± 0.01	0.31 ± 0.17	0.06 ± 0.02	0.02 ± 0.01	0.01 ± 0.01
Estate 26	26	1977-2004	576.0 ± 25.2	0.13 ± 0.03	0.02 ± 0.00	0.32 ± 0.05	0.04 ± 0.01	0.02 ± 0.00	0.00 ± 0.00
Estate 27	6	1994-1999	422.0 ± 0.0	0.18 ± 0.05	0.00 ± 0.00	1.16 ± 0.06	0.07 ± 0.01	0.03 ± 0.01	0.18 ± 0.06
Estate 28	12	1986-1997	224.5 ± 12.8	0.36 ± 0.07	0.03 ± 0.01	0.85 ± 0.05	0.11 ± 0.02	0.01 ± 0.00	0.04 ± 0.03
Estate 29	20	1977-2004	351.9 ± 11.9	0.26 ± 0.05	0.11 ± 0.01	0.51 ± 0.04	0.17 ± 0.02	0.07 ± 0.01	0.15 ± 0.07
Estate 30	11	1992-2003	458.5 ± 38.9	0.38 ± 0.05	0.04 ± 0.01	0.87 ± 0.08	0.11 ± 0.02	0.01 ± 0.00	0.02 ± 0.01
Estate 31	9	1996-2004	454.0 ± 17.3	1.54 ± 0.18	0.06 ± 0.02	0.26 ± 0.03	0.02 ± 0.00	0.01 ± 0.00	0
Estate 32	9	1993-2001	424.1 ± 12.2	0.16 ± 0.05	0.12 ± 0.01	1.14 ± 0.12	0.16 ± 0.03	0.37 ± 0.07	0.02 ± 0.00
Estate 33	9	1996-2004	317.1 ± 30.0	0.15 ± 0.03	0.05 ± 0.01	1.37 ± 0.23	0.08 ± 0.02	0.18 ± 0.04	0.04 ± 0.02
Estate 34	3	2002-2004	233.3 ± 31.7	0.71 ± 0.33	0.04 ± 0.02	0.97 ± 0.20	0.08 ± 0.04	0.01 ± 0.01	0.03 ± 0.01
Estate 35	3	2002-2004	288.0 ± 6.9	0.07 ± 0.05	0.03 ± 0.01	0.72 ± 0.39	0.04 ± 0.02	0.00 ± 0.00	0.00 ± 0.00
Estate 36	14	1988-2004	155.6 ± 12.5	0.14 ± 0.03	0.02 ± 0.00	1.00 ± 0.23	0.07 ± 0.02	0.00 ± 0.00	0.12 ± 0.04
Estate 37	3	2002-2004	224.0 ± 2.3	0.16 ± 0.01	0.09 ± 0.02	0.25 ± 0.02	0.04 ± 0.01	0.00 ± 0.00	0
Estate 38	14	1988-2003	122.7 ± 17.6	0.11 ± 0.03	0.02 ± 0.01	0.27 ± 0.04	0.07 ± 0.02	0.04 ± 0.01	0.00 ± 0.00
Estate 39	26	1979-2004	243.2 ± 16.2	0.72 ± 0.08	0.07 ± 0.01	0.85 ± 0.08	0.22 ± 0.02	0.34 ± 0.04	0.00 ± 0.00
Estate 40	24	1981-2004	68.7 ± 4.5	0.33 ± 0.05	0.01 ± 0.00	0.23 ± 0.03	0.07 ± 0.02	0.00 ± 0.00	0.00 ± 0.00
Estate 41	3	1996-1998	388.0 ± 74.2	0.07 ± 0.02	0.03 ± 0.01	0.69 ± 0.15	0.09 ± 0.08	0.00 ± 0.00	0.31 ± 0.07
Estate 42	5	1997-2004	405.0 ± 27.0	1.96 ± 0.76	0.18 ± 0.07	1.69 ± 0.46	0.25 ± 0.08	0.05 ± 0.02	0.02 ± 0.01
Estate 43	31	1973-2004	277.2 ± 5.5	0.12 ± 0.01	0.07 ± 0.01	0.10 ± 0.01	0.04 ± 0.01	0.01 ± 0.00	0.00 ± 0.00
Estate 44	7	1998-2004	116.7 ± 3.4	0.36 ± 0.10	0.11 ± 0.03	0.61 ± 0.08	0.06 ± 0.03	0.05 ± 0.02	0.36 ± 0.13
Estate 45	6	1998-2004	63.0 ± 7.3	4.77 ± 1.73	0.83 ± 0.20	2.32 ± 0.69	1.04 ± 0.26	0.12 ± 0.06	0.01 ± 0.01